



AD-A081 166 DEFENCE RESEARCH ESTABLISHMENT SUFFIELD RALSTON (ALBERTA) F/G 15/2  
EXCRETION OF INJECTED DIMETHYL MORPHOLINOPHOSPHORAMIDATE ON THE--ETC(U)  
JAN 80 M L MCDONALD  
UNCLASSIFIED DRES-TP-517 NL

| OF |  
AD  
A081166





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

ADA 081166

**DRES**

3

UNCLASSIFIED

UNLIMITED  
DISTRIBUTION

# SUFFIELD TECHNICAL PAPER

NO. 517

EXCRETION OF INJECTED DIMETHYL MORPHOLINOPHOSPHORAMIDATE  
ON THE BREATH OF RATS (U)

by

M.L. McDonald

PCN 13E30

January 1980

DTIC  
ELECTE  
FEB 27 1980

DDC FILE COPY



DEFENCE RESEARCH ESTABLISHMENT SUFFIELD : RALSTON : ALBERTA

**WARNING**

The use of this information is permitted subject to recognition  
of proprietary and patent rights .

80 2 26 085

UNCLASSIFIED

UNLIMITED  
DISTRIBUTION

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD  
RALSTON ALBERTA

⑨  
SUFFIELD TECHNICAL PAPER 517

⑭  
DRES-MP-

⑥  
EXCRETION OF INJECTED DIMETHYL MORPHOLINOPHOSPHORAMIDATE  
ON THE BREATH OF RATS (U)

by

⑩  
M.L. McDonald

⑪ Jan 80

⑫ 19

PCN 13E30

**WARNING**

The use of this information is permitted subject to recognition  
of proprietary and patent rights.

403104

UNCLASSIFIED

13

UNCLASSIFIED

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD  
RALSTON ALBERTA

SUFFIELD TECHNICAL PAPER NO. 517

EXCRETION OF INJECTED DIMETHYL MORPHOLINOPHOSPHORAMIDATE  
ON THE BREATH OF RATS (U)

by

M.L. McDonald

ABSTRACT

Male rats were injected with undiluted dimethylmorpholinophosphoramidate (DMMPA) into the peritoneal cavity. Levels on the exhaled breath were determined over the next 48 hours. Recovery was between 1% and 2% of the administered dose. The analysis indicated the presence of metabolites of the parent compound.

(U)

UNCLASSIFIED



# ACKNOWLEDGEMENT

The author wishes to thank the Chemistry Section, DRES, for carrying out the GLC analyses that were required during this study.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or special
PI	

UNCLASSIFIED

DEFENCE RESEARCH ESTABLISHMENT SUFFIELD  
RALSTON ALBERTA

SUFFIELD TECHNICAL PAPER NO. 517

EXCRETION OF INJECTED DIMETHYL MORPHOLINOPHOSPHORAMIDATE  
ON THE BREATH OF RATS (U)

by

M.L. McDonald

INTRODUCTION

Dimethyl morpholinophosphoramidate (DMMPA) has been used as a nerve agent simulant to assess standard operating procedures in chemical defence (1). Some of the work carried out to establish the safety of DMMPA has been concerned with the recovery of the simulant when it was applied to man and animals. Early studies (1,2) indicated that only 25 to 50% of the DMMPA applied to the skin of man could be found in the urine. Studies by Shaw and McDonald (3) showed that DMMPA was not accumulated in any of ten different rat tissues and organs. The maximum level in the tissues and organs was reached in about 18 hours and returned to near zero values within 48 to 72 hours. A balance study indicated that the major route of excretion was the urine, with small amounts appearing in the feces. Only 50% of the injected dose was recovered and the results showed no apparent breakdown of the DMMPA. Dawson et al. (4) investigating

UNCLASSIFIED

the persistence of DMMPA in various tissues of the rat also showed that significant amounts were found in the liver, kidneys and brain 4 hours after intraperitoneal (i.p.) injection and only trace amounts after 48 hours. They further reported that 58% of the administered dose was recovered in the urine, with the majority being excreted in the first 24 hours. It was stated that the DMMPA was probably metabolized in the body, however degradation could not be demonstrated in human plasma. Shaw and McDonald (3), based on observations made by Coleman (5), suggested that excretion on the breath probably accounted for the unrecovered DMMPA.

This study was carried out to establish whether or not DMMPA was excreted on the breath. A technique based on the miniature detector tube procedure for sampling DMMPA in the field (6) was developed to sample the exhaled breath of rats for DMMPA.

#### METHODS

A metabolic chamber was designed and built to provide a means of sampling the breath exhaled by the rat. The chamber was cylindrical in configuration and was made from a borosilicate glass cylinder of 100 mm ID with a total length of 37 cm (Fig. 1).

The design incorporated the use of two stainless steel screens of 3/8 inch and 1/8 inch mesh, respectively, positioned 25 mm apart (Fig. 2) and located at the bottom end of the chamber as shown in Figure 1. The upper screen, of 3/8 inch mesh, was the platform on which the rat remained during the sampling period. The urine and feces passed through to the bottom screen which retained the feces. The urine passed through to the bottom of the chamber to an outlet tube. To minimize any interruption of the chamber environment, a syringe connected to the outlet tube was used to withdraw the urine as required. It should be noted that the main purpose of the screen assembly was to insure maximum comfort for the animal.

An airflow ingress tube was located approximately 45 mm from the bottom of the chamber and 3 to 5 mm below the upper screen (Fig. 1).



A metered airflow was introduced through this tube over the rat and exited at the top of the chamber.

The air, containing the exhaled breath of the rat, flowed from the chamber via an 18 mm OD glass tube mounted in the centre of a No. 12 size rubber stopper. Twelve miniature detector tubes also fitted in the stopper in a circular configuration around the exit tube (Fig. 3) sequentially sampled the airstream prior to its exiting the chamber. The sampling procedure employed a second pump, a 12-position rotary valve with a crystal controlled electronic timer and miniature detector tubes charged with Tenax GC 60/80 mesh. The latter three items were supplied by the Chemistry Section, DRES, and are normally used for air sampling in the field. The method of connecting the miniature tubes to the rotary valve was modified to meet the requirements of the study (Fig. 4). The pump drew a metered flow of the chamber air to be sampled through a detector tube. A critical orifice monitored by a flow rate meter regulated the flow.

The activation of a tube from static to sampling mode was controlled by the rotary valve which was indexed through its 12 positions by the electronic timer. The timer was set for 1 hour sampling intervals per miniature detector tube.

Figure 5 illustrates the metabolic chamber with ancillary equipment.

Flow-rated miniature detector tubes supplied by the Chemistry Section were set up as follows. Approximately 40 mg of Tenax GC 60/80 mesh (Chromatographic Specialties Ltd.) was packed in a 38 mm length of borosilicate glass tubing of 4 mm OD and 2 mm ID. The Tenax GC bed was supported in the tube by stainless steel 200 mesh screens at each end. The tubes were conditioned for use by thermal desorption as per Chemistry Section's procedures. (The procedure is as follows: the tubes are placed in uncapped vials in quantities of 20 - 24; the vials are placed in a desiccator which is heated to 180°C in an oven and evacuated to  $5 \times 10^{-3}$  torr for 36 hours; the oven is then cooled, the vacuum pump turned off and the desiccator returned to atmospheric pressure by bleeding in super

pure nitrogen. Finally, the vials are capped and stored till the miniature tubes are required for sampling.)

The DMMPA was certified 99% pure as supplied from TeroChem Laboratories, Edmonton. It was stored at -66°C until received from the Chemistry Section for this study.

Male albino rats obtained from the University of Alberta (Sprague-Dawley), weighing between 190 and 250 grams, were used. Male and female rats, as reported by Coleman (5), respond identically to DMMPA. The rats were starved for 24 hours immediately prior to the run. This included control animals as well as those treated with DMMPA. At the end of the 24 hour starvation period the rats were injected (i.p.) with undiluted DMMPA. The doses used throughout the study were 0.1 mL (122.3 mg), 0.2 mL (245.6 mg) and 0.4 mL (489 mg). The weight of each dose level was calculated using a density of 1.2228 for DMMPA at 25°C. The injection site was swabbed with distilled water using a surgical gauze sponge to insure no surface DMMPA remained. The rat was then placed in the chamber and the chamber environment was sampled for DMMPA over the next 48 hours. The animal remained in the starved state while in the chamber. Control runs were carried out by placing untreated rats in the chamber immediately after 24 hours starvation and sampling the chamber environment for 48 hours.

The operating conditions for the metabolic chamber were as follows:

- flow rate through chamber = 1.0 L/min
- flow rate through miniature detector tubes while in sampling mode = 0.06 L/min average (a check made on this flow rate throughout a 48 hour run showed less than 5% variation)
- sampling time per miniature detector tube = 60 minutes
- temperature = ambient (air conditioned room)

The chamber was washed down between runs by submerging in chromic acid solution for 4 hours, then in 5% Contrad 70 solution overnight. It was then rinsed and dried at 75°C for 4 hours in a drying oven.

The miniature detector tubes (48 for a 48 hour run employing a 60 minute sampling period) were analyzed for DMMPA content via Gas Liquid

Chromatography (GLC) and the total recovery was calculated. The conditions employed for operation of the gas chromatograph were as follows:

- Instrument: Varian 1800 (data collection and processing carried out on a PDP 11/34)
- Detector: Flame photometric with phosphorus filter
- Column: Pyrex 6' x 1/8"
- Packing Support: Chromosorb W, Acid washed, DMCS treated, 80/100 mesh
- Stationary Phase: 2% Triton X305
- Column Temperature: 185°C
- Detector Temperature: 220°C
- Inlet Temperature: 230°C
- Carrier Gas: Nitrogen (Super Pure), flow rate: 35 mL/min @ 42 psi
- Inlet System: 3-way valve of local design (W.J. Fenrick)

The miniature detector tube was injected into the 3-way valve, allowed to heat for 20 seconds then flushed with nitrogen. Standardization was carried out by injecting a series of standards ranging from 3.3 ng to 6,600 ng DMMPA.

#### CALCULATIONS

Part I: Total mg/L DMMPA vapour concentration in miniature detector tubes:

$$S_{TV} = T_S \cdot F_1$$

$$X = \frac{X_1}{S_{TV}}$$

X = total mg/L DMMPA vapour concentration in miniature detector tubes

X<sub>1</sub> = the sum of the DMMPA (mg) recovered from the miniature



detector tubes (Normally this involved 48 tubes for a 48 hour run.)

$S_{TV}$  = total volume of chamber air sampled through the miniature detector tubes

$T_S$  = total sampling time in minutes (Normally 48 hours x 60 minutes/hour = 2880 minutes.)

$F_1$  = flow rate through miniature detector tubes in L/minute

Part II: Total DMMPA (mg) recovery:

$$R_T = A_{TV} \cdot X$$

$R_T$  = total DMMPA (mg) recovery

$A_{TV}$  = actual total volume (L) of sample stream that passed through chamber during the sampling period (Normally 1.0 L/min x 2880 minutes = 2880 L.)

#### RESULTS AND DISCUSSION

GLC analysis of the miniature detector tubes showed a DMMPA recovery of 1% to 2.2% of the total administered dose (Table I).

In addition to the DMMPA peak, the chromatograms of the breath from the 9 treated animals showed 2 and sometimes 3 small peaks. These were of a phosphorus bearing origin with a retention time less than DMMPA. These peaks were not present on the breath of the 4 untreated rats. The importance of this observation was the possibility that, metabolites of the parent compound were present. These peaks cannot be quantitated nor identified. It should be noted that due to their low magnitude, they would not add appreciably to the total recovery figure. Therefore, they were not taken into consideration for this purpose.

A recovery study was carried out to determine if any significant amount of DMMPA was adsorbed on the interior surfaces of the chamber, thus resulting in a low total recovery for the treated rat runs. Weighed amounts of DMMPA were introduced into the chamber and the sampling procedure for the vapour was run for 48 hours. The DMMPA was reweighed



and the miniature detector tubes were analysed and the percent recovery calculated. Several runs were carried out. These showed variable recovery, ranging from 82% to 95%. It was concluded that there was some loss due to adsorption on the interior surfaces in the chamber, but not of sufficient quantity to significantly increase the total recovery figure for the rat runs.

In answer to the original question, the study has indicated that rats injected (i.p.) with DMMPA do excrete it on the breath as well as possible metabolites of the parent compound. However, it is not excreted in sufficient quantity to explain the unaccounted for DMMPA (nearly 50%) in previous tissue, blood, feces and urine studies.

To fully determine the fate of injected DMMPA in the rat, further studies would have to be carried out. The most likely approach would be to confirm whether DMMPA is, or is not, metabolized in the body. To answer this question, a similar study to Shaw and McDonald (3) could be carried out using double-labelled DMMPA, with  $C^{14}$  in the methyl groups and tritium in the morpholine ring (3).

REFERENCES

1. McNally, W.D. and Adie, P.A. "Studies on the Total Intake Simulant DMMPA (U)". Suffield Technical Paper No. 469. 1977.  
UNCLASSIFIED. (Unclassified version of STP 410 originally issued August 1973.)
2. McNally, W.D. and Adie, P.A. "Urinary Excretion of DMMPA Following Its Inhalation (U)". Suffield Technical Note No. 267. 1973.  
UNCLASSIFIED.
3. Shaw, R.K. and McDonald, M.L. "Excretion of Injected Dimethylmorpholinophosphoramidate by the Female Albino Rat (U)". Suffield Technical Paper No. 479. 1977. UNCLASSIFIED.
4. Dawson, R.M., Lakeland, B.R., Poretski, M. and Bladen, M.P. "Dimethylmorpholinophosphoramidate (DMMPA) Persistence in Various Tissues of the Rat". Report MRL-R-705. January, 1978. Dept. of Defence, Materials Research Labs., Ascot Vale, Victoria, Australia.
5. Coleman, I.W. Private communication.
6. Pannekoek, W.J. Unpublished data.
7. Coleman, I.W. "The Toxicology of DMMPA Part I: The Acute Effects of DMMPA (U)". Suffield Technical Paper No. 466. 1977.  
UNCLASSIFIED.
8. Shaw, R.K. Private communication.

UNCLASSIFIED


TABLE I

DMMPA RECOVERY FROM THE BREATH OF THE RAT  
AFTER INTRAPERITONEAL INJECTION

Run	DMMPA Injected (mg)	Total Recovery (mg) Miniature Tubes (GLC Analysis)	% Recovery
1	125.5 (0.1 mL)	1.9270	1.57
2	125.5 (0.1 mL)	1.2289	1.00
3	125.5 (0.1 mL)	1.3599	1.11
4	245.0 (0.2 mL)	2.5816	1.05
5	245.0 (0.2 mL)	2.5761	1.05
6	245.0 (0.2 mL)	3.5614	1.45
7	490.0 (0.4 mL)	10.3130	2.10
8	490.0 (0.4 mL)	5.8539	1.19
9	490.0 (0.4 mL)	10.9300	2.23


UNCLASSIFIED






**FIG. 1 METABOLIC CHAMBER**






**FIG. 2: STAINLESS STEEL SCREEN ASSEMBLY**



**FIG. 3: MINIATURE DETECTOR TUBES  
AND CHAMBER EXIT TUBE**



**FIG. 4: ROTARY VALVE ASSEMBLY**




FIG. 5: METABOLIC CHAMBER WITH ANCILLARY EQUIPMENT



UNCLASSIFIED

Security Classification

**DOCUMENT CONTROL DATA - R & D**

(Security classification of title, body of abstract and indexing annotation must be entered when the overall document is classified)

1. ORIGINATING ACTIVITY  DEFENCE RESEARCH ESTABLISHMENT SUFFIELD		2a. DOCUMENT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>	
		2b. GROUP	
3. DOCUMENT TITLE  EXCRETION OF INJECTED DIMETHYL MORPHOLINOPHOSPHORAMIDATE ON THE BREATH OF RATS (U)			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		Technical Paper	
5. AUTHOR(S) (Last name, first name, middle initial)  McDonald, M.L.			
6. DOCUMENT DATE  JANUARY 1980		7a. TOTAL NO. OF PAGES  15	7b. NO. OF REFS  8
8a. PROJECT OR GRANT NO.  PCN 13E30		9a. ORIGINATOR'S DOCUMENT NUMBER(S)  SUFFIELD TECHNICAL PAPER NO. 517	
8b. CONTRACT NO.		9b. OTHER DOCUMENT NO.(S) (Any other numbers that may be assigned this document)	
10. DISTRIBUTION STATEMENT  UNLIMITED DISTRIBUTION			
11. SUPPLEMENTARY NOTES		12. SPONSORING ACTIVITY	
13. ABSTRACT  <p>Male rats were injected with undiluted dimethylmorpholinophosphoramidate (DMMPA) into the peritoneal cavity. Levels on the exhaled breath were determined over the next 48 hours. Recovery was between 1% and 2% of the administered dose. The analysis indicated the presence of metabolites of the parent compound.</p> <p style="text-align: right;">(U)</p>			



## KEY WORDS

DMMPA

Excretion

Breath

Rat

## INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the organization issuing the document.
- 2a. **DOCUMENT SECURITY CLASSIFICATION:** Enter the overall security classification of the document including special warning terms whenever applicable.
- 2b. **GROUP:** Enter security reclassification group number. The three groups are defined in Appendix 'M' of the DRB Security Regulations.
3. **DOCUMENT TITLE:** Enter the complete document title in all capital letters. Titles in all cases should be unclassified. If a sufficiently descriptive title cannot be selected without classification, show title classification with the usual one-capital-letter abbreviation in parentheses immediately following the title.
4. **DESCRIPTIVE NOTES:** Enter the category of document, e.g. technical report, technical note or technical letter. If appropriate, enter the type of document, e.g. interim, progress, summary, annual or final. Give the inclusive dates when a specific reporting period is covered.
5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the document. Enter last name, first name, middle initial. If military, show rank. The name of the principal author is an absolute minimum requirement.
6. **DOCUMENT DATE:** Enter the date (month, year) of Establishment approval for publication of the document.
- 7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.
- 7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the document.
- 8a. **PROJECT OR GRANT NUMBER:** If appropriate, enter the applicable research and development project or grant number under which the document was written.
- 8b. **CONTRACT NUMBER:** If appropriate, enter the applicable number under which the document was written.
- 9a. **ORIGINATOR'S DOCUMENT NUMBER(S):** Enter the official document number by which the document will be identified and controlled by the originating activity. This number must be unique to this document.
- 9b. **OTHER DOCUMENT NUMBER(S):** If the document has been assigned any other document numbers (either by the originator or by the sponsor), also enter this number(s).
10. **DISTRIBUTION STATEMENT:** Enter any limitations on further dissemination of the document, other than those imposed by security classification, using standard statements such as:
  - (1) "Qualified requesters may obtain copies of this document from their defence documentation center."
  - (2) "Announcement and dissemination of this document is not authorized without prior approval from originating activity."
11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.
12. **SPONSORING ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring the research and development. Include address.
13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document, even though it may also appear elsewhere in the body of the document itself. It is highly desirable that the abstract of classified documents be unclassified. Each paragraph of the abstract shall end with an indication of the security classification of the information in the paragraph (unless the document itself is unclassified) represented as (TS), (S), (C), (R), or (U).  
  
The length of the abstract should be limited to 20 single-spaced standard typewritten lines; 7 1/4 inches long.
14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a document and could be helpful in cataloging the document. Key words should be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context.